



RESEARCH ARTICLE - ANTS

Richness of Termites and Ants in the State of Rio Grande do Sul, Southern Brazil

E DIEHL¹, E DIEHL-FLEIG¹, EZ DE ALBUQUERQUE², LK JUNQUEIRA³1 - *Universidade Federal do Rio Grande do Norte, Natal, RN, Brazil.*2 - *Museu de Zoologia, Universidade de São Paulo, São Paulo, SP, Brazil.*3 - *Pontifícia Universidade Católica de Campinas, Campinas, SP, Brazil.*

Article History

Edited by

Gilberto M M Santos, UEFS, Brazil

Received 18 December 2013

Initial acceptance 28 January 2014

Final acceptance 06 April 2014

Keywords

Social insects, distribution, altitude.

Corresponding author

Eduardo Diehl-Fleig

Instituto do Cérebro, Universidade

Federal do Rio Grande do Norte

Av. Nascimento de Castro 2155

Natal, RN, Brazil

59056-450

E-mail: edf.formiga@gmail.com

Abstract

Previous studies on the effects of environmental factors, such as altitude, latitude, temperature, deforestation, forest fragmentation, fire, and flood on the community structure of termites and ants were conducted in various regions of Brazil; few of them were carried out in the southernmost Brazilian state of Rio Grande do Sul. Here we describe termites and ants diversity at different sites along the four geomorphologic units of this state. We recorded 16 taxa of termites, of which three are new state records, increasing to 19 the number of termite species known to occur in the state. Accordingly, we also found 73 species and 115 morphospecies of ants, of which only one was a new record, raising to 265 taxa the number of ant species known to occur in the state. As expected, we found a higher species richness of ants than termites. The low richness of both groups relative to other Brazilian regions could be a consequence of the subtropical to temperate climate in the state, since most portions of the state are below 30° latitude, the study areas be above 500 m altitude, and other environmental characteristics of each site. We suggest a positive relationship between species richness of termites and altitude, while ant richness indicated an inverse relationship. However, our data are not conclusive, due to the low number of replications in each altitude, particularly for termites. This study is unique in presenting an updated checklist of termites and ants in the state of Rio Grande do Sul.

Introduction

Studies on biodiversity are urging in face of diversity loss (Wilson, 1997; McGeoch & Chown, 1998). Termites (Mill, 1982; Constantino, 1992; Eggleton et al., 1995; Black & Okwakol, 1997; Lavelle et al., 1997; Jones, 2000; Gathorne-Hardy et al., 2001; Jones et al., 2003) and ants (Majer, 1983; Hölldobler & Wilson, 1990; Andersen, 1997; Silva & Brandão, 1999; Ward, 2000; Ribas et al., 2012) have been considered biodiversity indicators due to their biological and ecological characteristics. These groups have also been considered as surrogate groups in evaluations of conservation status, degradation, or recovery of terrestrial ecosystems (New, 1996; Majer, 1996; Andersen, 1997; Majer & Nichols, 1998; Lobry de Bruyn, 1999; Jones & Eggleton, 2000; Bandeira &

Vasconcello, 2002; Diehl et al., 2004).

Like ants, termites are entirely eusocial and have profound ecological significance in the tropics (Engel et al., 2009). Studies have shown that ants and termites play a crucial role to create soil structure, influence aeration, water infiltration and nutrient cycling acting as ecosystem engineers (Lobry de Bruyn & Conacher, 1990; Lavelle et al., 1997; Mora et al., 2005; Jouqueta et al., 2011; Del Toro et al., 2012). In some cases, termites and ants are also responsible for increased crop yield under dry conditions through soil water infiltration due to their tunnels and improved soil nitrogen (Evans et al., 2011).

Environmental disturbances, such as deforestation, forest fragmentation, fire, and floods may affect the community structure of termites (De Souza & Brown, 1994; Eggleton et



al., 1994, 1995; Davies, 2002; Bandeira et al., 2003; Inoue et al., 2006) and ants (Hölldobler & Wilson, 1990; Majer & Nichols, 1998; Diehl et al., 2004; Schmidt & Diehl, 2008). Previous studies have shown that local species richness of ants and termites is influenced by environmental factors, such as temperature, rainfall, and vegetation (Benson & Harada, 1988; Silva & Brandão, 1999; Gathorne-Hardy et al., 2001; Oliveira & Del-Claro, 2005; Del-Claro, 2008; Jones & Eggleton, 2011). There has also been some evidence that increasing altitude and decrease of temperature decrease species richness (Kusnezov, 1957; Collins, 1980; Ward, 2000; Gathorne-Hardy et al., 2001) and that the phylogenetic structure might be involved in this process, specially in the temperate forests (Machac et al., 2011).

The Neotropics rank in the third position in termite richness among the biogeographic regions, with the Oriental and Ethiopian ones presenting the highest number of species recorded so far (Krishna et al., 2013). About 72% of the neotropical species occur in South America (Constantino, 2014) and the sites with good termite surveys are between the parallels 0° S and 20° S, which include the biomes Amazon and Atlantic Forests, Cerrado and Caatinga (Constantino & Acioli, 2006). A review of the termite fauna in across Brazil can be found in Constantino & Acioli (2006). Recent studies recorded 33 species for five fragments of semideciduous Atlantic Forest in northern Brazil, varying locally from 11 to 27 species (Souza et al., 2012). With different levels of perturbation, 26 species were recorded in a region of semi-arid Caatinga (Vasconcellos et al., 2010). In southern Brazil, termite fauna is poorly known. Studies mention only 16 termite species to Rio Grande do Sul (Araújo, 1977; Constantino, 1998; Fontes, 1998; Castro & Diehl, 2003; Diehl et al., 2005b; Florencio & Diehl, 2006). This low number can be due to the climate (Eggleton et al., 1994; Constantino & Acioli, 2006), or lack of taxonomists and local studies (Diehl-Fleig et al., 1995; Constantino & Acioli, 2006).

Despite its importance, little is known about the edaphic insects, specially ants and termites, in southern Brazil. Thus, here we present the richness and a checklist of termites and ants in the four geomorphic units of Rio Grande do Sul, reporting new records for state.

The checklist also includes species from other inventories conducted in the state of Rio Grande do Sul by researchers apart from our group led by E. Diehl. Most aimed at collecting species of *Acromyrmex* and *Atta* or urban ant fauna and few have focused on ant species inventories in forests and other relevant ecosystems (please see supplementary material for further details at):

<http://periodicos.uefs.br/ojs/index.php/sociobiology/rt/suppFiles/313/408>

DOI: 10.13102/sociobiology.v61i2.145-154.s451

Material and Methods

Study areas

Geomorphological units of Rio Grande do Sul

The southernmost Brazilian state of Rio Grande do Sul has four geomorphologic units: Southern Plateau, Central Depression, Sul-Riograndense Shield, and Coastal Plain (Fig 1). Two main vegetation types originally occurred in the state: forests and grasslands occupying 34% and 46% of the state area, respectively. The remaining was occupied by coastal vegetation, wetlands, and other vegetation types. The vegetation reflects soil type and origin and forests usually occur where the soil is deep and fertile. Grasslands develop where the soil is siliceous or shallow. While the southern region of the state is dominated by grasslands, the northern grasslands are interspersed with Araucaria forest (Rambo, 1994).

The weather in southern Brazil is subtropical humid, according to the Köppen classification. While the average annual temperature in the Coastal Plain is around 18°C, temperature approaches 20°C in the Central Depression, which also has mild winters. Conversely, the annual average temperature in the Southern Plateau does not exceed 15°C, due to its high altitude, winters are harsh, with often negative temperatures, severe frosts, and even deep snow. The Campanha, in the Sul-Riograndense Shield, is an open landscape and receives all continental winds in the winter, but due to their large grassland areas it receives much sunlight resulting in summers with high temperatures (Rambo, 1994).

Sampling sites (Fig 1) are described below and are mostly under different levels of disturbance:

- 1) remnants of seasonal semideciduous forest and suburban areas around São Leopoldo (29° 45' - 29° 47' S, 51° 05' - 51° 10' W; 15 m a.s.l.) in the Central Depression (Mayhê-Nunes & Diehl-Fleig, 1994; Diehl-Fleig & Fleig, 1997; Haubert et al., 1998; Flores et al., 2002; Diehl et al. 2006; Florencio & Diehl, 2006; Marchioretto & Diehl 2006);
- 2) periurban and urban areas in Canela (29° 21' S, 50° 49' W; 819 m a.s.l.) and Gramado (29° 22' S, 50° 52' W; 825 m a.s.l.) in the Southern Plateau (Diehl-Fleig, 1997);
- 3) dunes and restingas in Torres (29° 20' - 29° 23' S, 49° 43' - 49° 46' W; 0 - 16 m a.s.l.) in the northern Coastal Plain (Diehl-Fleig et al., 2000; Hameister et al., 2003; Albuquerque et al. 2005; Diehl, unpub. ms.);
- 4) grasslands in the Caçapava do Sul (30° 47' S, 52° 24' W; 220 m a.s.l.) in soils with different copper levels, including a copper mine and a native savanna area in the Camaquã River Basin in the Sul-Riograndense Shield (Diehl et al., 2004);
- 5) restingas in Morro da Grota (30° 21' S, 50° 01' W; 263 m a.s.l.) near Lagoa dos Patos and a typical forest of granite

- soil, sandy and rocky areas in Pedreira beach (30° 21' S, 50° 02' W; at the sea level), on the banks of the Guaíba Lake, both at the Itapuã State Park in Viamão, which is within both Sul-Riograndense Shield and Coastal Plain (Sacchett & Diehl, 2004; Diehl et al., 2005a; Diehl, unpub. ms.);
- 6) Eucalyptus plantation in restingas in Capivari do Sul (30° 11' S, 50° 21' W; 12 m a.s.l.) and Tramandai (29° 59' S, 50° 13' W; 8 m a.s.l.) in the northern Coastal Plain (Fonseca & Diehl, 2004);
 - 7) wetland used as pastures for cattle or irrigated rice in Santo Antônio da Patrulha (29° 54' S, 50° 33' W; 25 m a.s.l.) in the Coastal Plain (Diehl et al., 2005b; Moraes & Diehl, 2009);
 - 8) an area of mixed ombrophilous forest in the São Francisco de Paula National Forest (29° 23' - 29° 27' S, 50° 23' - 50° 25' W; 930 m a.s.l.) and in the Pró-Mata Research and Nature Conservation Center (29° 28' S, 50° 13' W; 900 a.s.l.) in the Southern Plateau (Diehl et al., 2005c; Pinheiro et al., 2010);
 - 9) vineyards (29° 56' S, 51° 33' W; 645 m a.s.l.) in Bento Gonçalves, within the northeastern Central Depression (Sacchett et al., 2009);
 - 10) yerba mate plantations in Mato Leitão (29° 31' S, 52° 07' W; 81 m a.s.l.), Ilópolis (28° 55' S, 52° 07' W; 683 m a.s.l.), and Putinga (29° 00' S, 52° 09' W; 435 m a.s.l.) in the northeastern Central Depression (Junqueira et al., 2001; Steffens, 2006);
 - 11) primary forest and reforestation area in Alto Ferrabraz, Sapiranga (29° 34' S, 50° 56' W; 500 m a.s.l.). This is a transition zone between the Southern Plateau and the northeastern Central Depression (Haubert, 2006);
 - 12) secondary forest, with Acacia plantation and an area with reforestation in Rolante (29° 36' 32.2" S, 50° 31' 39.1" W; 370 m a.s.l.) in the Central Depression (Schmidt & Diehl, 2008);
 - 13) grasslands in Campos de Cima da Serra (29° 10' S, 50° 10' W; 1,200 m a.s.l.) in the Southern Plateau (Albuquerque & Diehl, 2009);

At each termite sampling site, we established a transect (100 m x 3 m), further divided into 20 sections of 15 m². We then sampled ten non-contiguous sections on the right and left sides of the transect alternately, with a sampling effort of 60 minutes per section. In each section, we search for termites in leaf litter and humus at the base of trees and roots, under both rocks and fallen trees, inside logs, hollow twigs, decomposing fallen branches, epigeal and arboreal nests. We also set up one Termitrap[®] bait per section (Almeida & Alves, 1995) at 15 cm deep, 60 days before samplings. Furthermore, we also searched for termites in ten blocks (340 cm³) of soil horizon A per section.

Termites were stored in individual amber glass with 80% ethanol. Termites were identified to the genus level




Fig 1. Distribution of combined termite and ant sampling (crossed circles) and other ant inventoried sites (open circles) in the four geomorphologic units of Rio Grande do Sul, southern Brazil by Diehl group (geomorphologic map adapted from SEPLAG, 2008).

(Fontes, 1995; Constantino, 1999; 2002) and eventually to the species level or morphospecies. We also consulted specialist taxonomists to confirm the identification of termites.

We used five techniques to sample ants: 1) manual capture of all ants on the ground, under stones, in hollow trees, and branches; 2) baits with sardines in vegetable oil on a piece of filter paper (6 cm²); 3) leaf litter samples, firstly sieved in the field and further using Winkler extractors, in which they remained for 72 hours; 4) pitfall traps with and without attractive, which consisted of 200-300 mL plastic cups buried up to the top containing 70% ethanol, remaining on the ground for 24 to 48 hours. Attractive traps were composed by sardines in vegetable oil pierced in a galvanized wire rod; 5) underground trap with attractive, which was a small plastic pot (3.3 cm wide by 5 cm high) with small holes on the sides. Within each trap a piece of 1 cm³ sardines in vegetable oil was added. Traps were buried 20 cm deep by 48 hours. In grasslands we used direct sampling, sardine baits and pitfall traps. In dunes, we only used direct sampling and sardine baits. At each site, we established two to three 100 m transects, along which we sampled every 10 m. Sampling effort was at least 2 hours in each site.

We classified ants into subfamilies following Bolton (2014) and by genus following Bolton (1994) and Fernández (2003). The subfamilies Ponerinae and Ecitoninae were treated according to the most recent reviews - see supplementary material for details. We used several identification keys and also consulted specialists to identify species. We assigned specimens to morphospecies when specific identification

was not possible. At least three specimens of each species/morphospecies were mounted with entomological pins and the remaining preserved in 70% ethanol.

Specimens of termites and ants are housed at the Collection of Social Insects (Isoptera and Formicidae) of the Universidade do Vale do Rio dos Sinos (UNISINOS), São Leopoldo, Rio Grande do Sul, Brazil.

The checklist presented results of ca. 30 years of collections led by E. Diehl added to most ant inventories conducted by few other researchers in the state of Rio Grande do Sul. We reviewed papers published from 1972 to 2014 included in the databases ISI Web of Science, Scielo and Google Scholar that contained the keyword combination: ant OR Formicidae AND Rio Grande do Sul. Only papers referring nominal species and site location (municipality or more specific location) were considered, except those with doubtful identification (approximate identification, to be confirmed, compared to, of group, subgenus or species complex). Three studies on *Atta* and *Acromyrmex* distribution were excluded because site location was not provided. Species previously registered to the state was provided by the catalog of Kempf (1972) with additions of Brandão (1991).

Results

We recorded 16 isopterans from two families (Table 1). We only found one species of the Kalotermitidae genus *Rugitermes*. We also recorded a new genus of Termitidae, besides seven morphospecies and seven species from four subfamilies (Apicotermittinae, Nasutitermitinae, Syntermitinae, and Termitinae).

Termite species richness was very low. We recorded 10 species in the Southern Plateau (181-1,200 m a.s.l.) and 15 in the Central Depression (20-30 m), whereas in the Sul-Riograndense Shield (23-29 m) and northern Coastal Plain (5-15 m) we found the lowest richness: four and six species, respectively.

We found 26 ant genera from seven subfamilies (Dolichoderinae, Dorylinae, Ectatomminae, Formicinae, Heteroponerinae, Myrmicinae, and Ponerinae) that were stored in 70% ethanol. Furthermore, we recorded 51 genera from nine subfamilies (Amblyoponinae, Dolichoderinae, Dorylinae, Ectatomminae, Formicinae, Heteroponerinae, Myrmicinae, Ponerinae, and Pseudomyrmecinae), which were stored in entomological drawers. In total, we recorded 73 species and 115 morphospecies in Rio Grande do Sul. Compiling only nominal ant species registered by our group and other researchers, 127 species distributed among 55 genera from 10 subfamilies occur in the state (an extended list is available as supplementary material).

A single species, *Monomorium floricola*, is reported as new record for the state of Rio Grande do Sul. Thus, ant species richness in Rio Grande do Sul raises to 265, considering published data on nominal species from the catalogs (Kempf, 1972; Brandão, 1991), from our research group and the other groups. This number would likely increase, if we

Table 1. Termites recorded in the state of Rio Grande do Sul, Brazil.

Family/ Subfamily	Taxon
Kalotermitidae	<i>Rugitermes</i> sp. (1, 3, 4)*
Termitidae	
Apicotermittinae	Apicotermittinae 1 (1, 2, 3, 4) Apicotermittinae 2 (1, 2, 3, 4) <i>Grigiotermes bequaerti</i> (Snyder & Emerson, 1949) (3) <i>Grigiotermes</i> sp. (3) <i>Ruptitermes</i> sp. (3) <i>Tetimatermes</i> sp. (3) New Genus 1 (3)
Nasutitermitinae	<i>Araujotermes caissara</i> Fontes, 1982 (1) <i>Cortaritermes fulviceps</i> (Silvestri, 1901) (1, 2, 3, 4) <i>Nasutitermes aquilinus</i> (Holmgren, 1910) (3, 4) <i>Nasutitermes jaraguae</i> (Holmgren, 1910) (3, 4)
Syntermitinae	<i>Cornitermes cumulans</i> (Kollar, 1832) (2, 3, 4)
Termitinae	<i>Dihoplotermes inusitatus</i> Araújo, 1961 (3, 4) <i>Neocapritermes</i> sp. (1, 3, 4) <i>Termes</i> sp. (3, 4)

*Geomorphologic units: 1, Coastal Plain; 2, Sul-Riograndense Shield; 3, Central Depression; 4, Southern Plateau

consider the morphospecies not yet identified to the species level and other published studies with nominal species and site location indicated (see studies available as supplementary material).

Ant richness and composition varied along the geomorphologic units. We found 88 species/morphospecies in the Southern Plateau (900-1,200 m a.s.l.), 40 in the transition between the Southern Plateau and the Central Depression (500 m), 108 in the Central Depression (10-683 m), 51 in the Sul-Riograndense Shield (220 m), 81 in the transition between the Sul-Riograndense Shield and the Coastal Plain (16-263 m), and 97 in the Coastal Plain (0-80 m). Of the 188 species/morphospecies, most (40.43%) were restricted to a single geomorphologic unit, while only ca. 2.5% were common to all units (*Pheidole* sp.3, *Pheidole* sp.6, *Pheidole* sp.15, *Wasmannia* sp. and *Wasmannia* sp.1).

Excluding three sites where ant fauna was inventoried exclusively in agricultural or urban areas, there is some indication that species richness and composition could vary in response to an altitudinal gradient. The highest ant richness, 56, 81, and 73 species were found in sites at the sea level, 10 m, and 26 m, respectively. Conversely, we found 51, 35, 39, 55, and 33 species in sites at 220 m, 370 m, 500 m, 930 m, and 1,200 m, respectively. Of the 158 ant species/morphospecies in these eight areas, none were present in all sites, only ca. 3% were common to seven areas. The great majority (35.44 %) of ants were restricted to a single site.

Discussion

According to Krishna et al. (2013) there are 2,937 termite species worldwide, of which 569 are found in the Neotropics. The On-line Termite Database maintained by Constantino (2014) refers 2,882 termite species in the world, of which 562 species occur in the Neotropics (database was last updated in September 2012). Estimates of termite richness for Brazil are not yet accurate, ranging from 250 to 364 species (Canello & Schlemmermeyer, 1999; Constantino, 1999; 2014; Fontes & Araújo, 1999). However, the estimated species richness should increase with termite sampling in other areas of the country. Our results provide records of a new genus likely belonging to Apicotermittinae, another unidentified species of this subfamily (Apicotermittinae 1), and *Dihoplotermes inusitatus*. Thus, the number of known termite species in Rio Grande do Sul increases to 19.

Drywood termites of the family Kalotermitidae have major economic importance and a wide geographic distribution. These termites are poorly known because they inhabit places difficult to access (Jones & Eggleton, 2011). The largest and most diverse isopteran family is Termitidae (Eggleton & Tayasu, 2001). We found species from four out of the eight termite subfamilies previously known to occur in Rio Grande do Sul (Apicotermittinae, Nasutitermitinae, Syntermitinae, and Termitinae). A negative correlation between altitude and species richness has been reported for various organisms (Rahbek, 2005; McCain, 2009; McCain & Grytnes, 2010). This correlation seems to be dependent on altitude in tropical and subtropical areas. Altitude is positively correlated with richness in latitudes below 30° and below 500 m, but negative in latitudes above 30° and at altitudes above 500 m (Kusnezov, 1957; Ward, 2000).

We found indication that termite species richness relates positively to altitude. Previous studies (Inoue et al., 2006) found the species richness of Nasutitermitinae increased with altitude, while species richness and abundance of Macrotermittinae decreased with altitude. The other groups were apparently not influenced by altitude. Recently, Palin et al. (2010) evaluated the influence of the variation along an Amazon–Andes altitudinal gradient in Peru on richness, abundance, and diversity of functional groups of termites. They registered 49 species and verified that, in general, the diversity declined with increased elevation. The functional groups responded differently to the upper distribution limit: for the soil-feeding it was between 925 and 1,500 m a.s.l., while the wood-feeding termites was between 1,550 and 1,850 m a.s.l. And this differential response led the authors to suggest that the energy requirements for each group are a key factor in shaping their occurrence associated with the altitude and temperature.

Termite species richness seems to be influenced by local environmental conditions, such as altitude, temperature, rainfall, and vegetation (Collins, 1980; Gathorne-Hardy et al.,

2001; Davies, 2002; Inoue et al., 2006; Jones & Eggleton, 2011). The species richness of termites we found is as low as reported in subtropical and temperate areas. For example, previous studies found two to eight morphospecies in the Atlantic Forest of southeastern Brazil, whereas an average of 30 morphospecies were found in tropical forests near Bahia, northeastern Brazil (Canello et al., 2002).

The far southernmost part of South America has often been considered as a separate biogeographic sub-region. It is depauperate in termite species due to its high latitude (Eggleton, 2000), and countries like Chile and Uruguay have even fewer species registered: six and nine, respectively, while 37 are reported for Paraguay (Constantino, 2014). Argentina, on the other hand, has 57 species reported (Constantino, 2014), even though this number could increase to 95 (Torales et al., 2008). Argentina local termite species varies from 12 in the Chaco province (Godoy et al., 2012) to 26 species in north-east Argentina (Laffont et al., 2004).

We present the first two records of the rare, monotypic genus *Dihoplotermes* to Rio Grande do Sul. A colony of *D. inusitatus* was found inhabiting a nest *Cornitermes bequaerti* in gallery forests in Mato Grosso (Mathews, 1977), whereas other studies (Constantino, 1999) found *D. inusitatus* in the Cerrado and disturbed habitats in the Southeast. We found the first record of *D. inusitatus* within a nest of *C. cumulans* in grasslands surrounding the São Francisco de Paula National Forest. However, we could not find this species in further surveys. This is a rare species that could have went locally extinct after a *Pinus* plantation covered the area. The other record of *D. inusitatus* was also in a nest of *C. cumulans* in the suburb of São Leopoldo, more than 500 km away from the first site. With the increase of urbanization in the region, this rare species is subject to rapid local extinction.

The family Formicidae comprises more than 15,700 species and subspecies (AntWeb, 2013); recent estimates suggest that this number should exceed 23,000 species (AntWeb, 2013). However, previous studies mention 208 (Kempf, 1972) or 224 species Brandão (1991) to Rio Grande do Sul. Since its last catalog update (Brandão, 1991), 23 years ago, our study is the first to present an updated checklist of the ant species in Rio Grande do Sul. If we take into account the determination of species level of the all morphospecies in our collection at Unisinós and in other studies, the species list would likely increase. Only few list are available for other states, such as Santa Catarina (Ulysséa et al., 2011), where ant richness (366 species and 17 subspecies) is greater than the 265 species now updated to Rio Grande do Sul. Apart from different collection efforts, historical aspects linked to the myrmecology development are part of this numerical difference reported.

Ant inventories in Rio Grande do Sul until the early 2000s focused primarily in the genera *Atta* and *Acromyrmex* (e.g., Diehl-Fleig, 1997). This attines have long been known for their pest status and major impact on agriculture, fairly

common and economically relevant throughout the state. It is clear that ant fauna inventories in RS are still very scarce and the geomorphic units are undersampled, particularly the Pampa biome. This grassland ecosystem is unique in fauna and vegetation among Brazilian biomes and less than 42% remains preserved (Roesch et al., 2009). So far, ant fauna has been surveyed mostly in the Atlantic Forest biome in the state (Diehl et al., 2005c; Albuquerque & Diehl, 2009; Pinheiro et al., 2010) while a single study included the Pampa (Marques & Schoereder, 2014).

The diversity of habitats, vegetation, altitude, latitude, and climate between sampling sites should be the main factors responsible for the differences found in termite and ant species richness along the study areas and also other tropical areas in Brazil (Oliveira & Del-Claro, 2005). Currently, 106 ant genera are known to occur in Brazil (Antwiki, 2013), while the termite diversity is much lower, only 74 genera (Constantino, 2014). As expected, we found a higher richness of ant genera than termite along the studied areas. Species diversity decreases with increasing distance from the equator (Kusnezov, 1957; Eggleton et al., 1994; Ward, 2000). Therefore, the lowest species richness of termites and ants found by us was expected. Additionally, the low richness may be due to the subtropical climate, the low latitude (< 30°), and high altitudes (> 500 m), as might be case for our findings that ant richness was lower at higher altitudes. Moreover, sampled sites varied in the degree of habitat complexity, from single crop to native forest, which also play a key role in shaping species richness and composition with different outcomes (e.g., Lassau & Hochuli, 2004; Silva et al., 2007; Pacheco & Vasconcelos, 2012). This study is unique in compiling the richness of both termites and ants so far known in the state of Rio Grande do Sul and we hope it provides a landmark in exposing the necessity for future studies and establishing efforts in unraveling our eusocial insect diversity.

Acknowledgements

We would like to express our gratitude to three anonymous reviewers, to Dr. Jacques H. Delabie, Dr. Antônio J. Mayhé-Nunes, Dr. Rodrigo dos S.M. Feitosa, Dr. Carlos E. Sanhudo and Thiago Ranzani who helped us to confirm either/or identify ant species, in addition to providing several dichotomous keys. We also thank Dr. Reginaldo Constantino and Dr. Luiz R. Fontes for identification of termites. Daniela F. Florencio, Laura V.A. Menzel and Carlos E. Sanhudo helped in field and laboratory work. FAPESP provided financial support to E.Z. de Albuquerque (Proc. N. 2010/02560-5). E. Diehl was supported by CNPq. Formigas do Brasil research group provided much of the ant bibliographic review. We thank Unisinos for the laboratory facilities..

References

- Albuquerque, E.Z.; Diehl-Fleig, Ed. & Diehl, E. (2005). Density and distribution of nests of *Mycetophylax simplex* (Emery) (Hymenoptera: Formicidae) in areas with mobile dunes on the northern coast of Rio Grande do Sul, Brazil. *Revista Brasileira de Entomologia*, 49(1): 123-126. doi: 10.1590/S0085-56262005000100013
- Albuquerque, E.Z. de, Diehl, E. (2009). Análise faunística das formigas epígeas (Hymenoptera, Formicidae) em campo nativo no Planalto das Araucárias, Rio Grande do Sul. *Revista Brasileira de Entomologia*, 53: 123-126. doi: 10.1590/S0085-56262009000300014
- Almeida, J.E.M. & Alves, S.B. (1995). Seleção de armadilhas para captura de *Heterotermes tenuis* (Hagen). *Anais da Sociedade Entomologica do Brasil*, 24: 619-624.
- Andersen, A.N. (1997). Using ants as bioindicators: multi-scale issues in ant community ecology. *Conservation Ecology* 1: 8. Retrieved from: <http://www.consecol.org/vol1/iss1/art8>
- AntWeb. (2013). <http://www.antweb.org>. (accessed date: 8 December, 2013).
- Antwiki (2013). http://www.antwiki.org/wiki/Distribution_and_Diversity. (accessed date: 27 November, 2013).
- Araújo, R.L. (1977). *Catálogo dos Isoptera do Novo Mundo*. Rio de Janeiro: Academia Brasileira de Ciências, 92 p.
- Bandeira, A.G. & Vasconcellos, A. (2002). A quantitative survey of termites in a gradient of disturbed highland forest in Northeastern Brazil (Isoptera). *Sociobiology*, 39: 429-439.
- Bandeira, A.G., Vasconcellos, A., Silva, M.P. & Constantino, R. (2003). Effects of habitat disturbance on the termite fauna in a highland humid forest in the Caatinga domain, Brazil. *Sociobiology*, 42: 117-127.
- Benson, W.W. & Harada, A.Y. (1988). Local diversity of tropical and temperate ant faunas (Hymenoptera: Formicidae). *Acta Amazonica*, 18(3-4): 275-289.
- Black, H.I.J. & Okwakol, M.J.N. (1997). Agricultural intensification, soil biodiversity and agroecosystem function in the tropics: the role of termites. *Applied Soil Ecology*, 6: 37-53. doi: 10.1016/S0929-1393(96)00153-9
- Bolton, B. (1994). *Identification guide to the ant genera of the world*. Cambridge: Harvard University Press, 222 p
- Bolton, B. (2003). *Synopsis and classification of Formicidae*. Florida: *Memoirs of the American Entomological Institute*, 370 p
- Brandão, C.R.F. (1991). Adendos ao catálogo abreviado das formigas da região Neotropical (Hymenoptera: Formicidae). *Revista Brasileira de Entomologia*. 35: 319-412.
- Cancello, E.M. & Schlemmermeyer, T. (1999). *Reino Animalia*:

- Ordem Isoptera. In C.R.F. Brandão & E.M. Cancellato (Eds.), Biodiversidade do Estado de São Paulo, Brasil: síntese do conhecimento ao final do século XX, invertebrados terrestres (pp. 82-91). São Paulo: FAPESP.
- Cancellato, E.M.; Oliveira, L.C.M.; Reis, Y.T. & Vasconcelos, A. (2002). Termite diversity along the Brazilian Atlantic Forest. In Proceedings of the XIV International Congress of IUSI (pp. 164). Sapporo: Hokkaido Univ.
- Castro, Z. & Diehl, E. (2003). Gêneros de térmitas em ninhos epigeos no campus da Unisinos, São Leopoldo, RS. *Acta Biologica Leopoldensia*, 25: 93-102.
- Collins, N.M. (1980). The distribution of soil macrofauna on the west ridge of Gunung (Mount) Mulu, Sarawak. *Oecologia*, 44: 263-275. doi: 10.1007/BF00572689
- Constantino, R. (1992). Abundance and diversity of termites (Insecta: Isoptera) in two sites of primary rain forest in Brazilian Amazonia. *Biotropica*, 24: 420-430.
- Constantino, R. (1998). Catalog of the living termites of the new world (Insecta: Isoptera). *Arquivos de Zoologia*, 35: 135-260. doi: 10.11606/az.v35i2.12014
- Constantino, R. (1999). Chave ilustrada para identificação dos gêneros de cupins (Insecta: Isoptera) que ocorrem no Brasil. *Papéis Avulsos de Zoologia*, 40: 387-448.
- Constantino, R. (2002). An illustrated key to Neotropical termite genera (Insecta: Isoptera) based primarily on soldiers. *Zootaxa*, 67: 1-40.
- Constantino, R. (2014). On-line termite database. <http://164.41.140.9/catal/>. (accessed date: 10 February, 2014).
- Constantino, R. & Acioli, A.N.S. (2006). Termite diversity in Brazil (Insecta: Isoptera). In F.M.S. Moreira, J.O. Siqueira & L. Brussaard (Eds.), *Soil biodiversity in Amazonian and other Brazilian ecosystems*. (pp. 117-128). London: CAB International. doi: 10.1079/9781845930325.0117
- Davies, R.G. (2002). Feeding group responses of a Neotropical termite assemblage to rain forest fragmentation. *Oecologia*, 133: 233-242. doi: 10.1007/s00442-002-1011-8
- De Souza, O.F.F. & Brown, V.K. (1994). Effects of habitat fragmentation on Amazonian termite communities. *Journal of Tropical Ecology*, 10: 197-206. doi: 10.1017/S0266467400007847
- Del-Claro, K. (2008). Biodiversidade Interativa: a ecologia comportamental e de interações como base para o entendimento das redes tróficas que mantém a viabilidade das comunidades naturais. In J. Seixas & J. Cerasoli (Eds.), *UFU, ano 30 – Tropeçando Universos (artes, humanidades, ciências)* (pp. 599-614). Uberlândia: EDUFU.
- Del Toro, I., Ribbons, R.R. & Pelini, S.L. (2012). The little things that run the world revisited: a review of ant-mediated ecosystem services and disservices (Hymenoptera: Formicidae). *Myrmecological News*, 17: 133-146
- Diehl-Fleig, E., Silva, M.E. da & Castilhos-Fortes, R. (1995). O problema dos cupins no Rio Grande do Sul. In E. Berti Filho & L.R. Fontes (Eds.), *Alguns aspectos atuais da biologia e ecologia dos cupins* (pp. 53-56). Piracicaba: FEALQ.
- Diehl, E. (1997). Ocorrência de *Acromyrmex* em áreas com distintos níveis de perturbação antrópica no Rio Grande do Sul. *Acta Biologica Leopoldensia*, 19: 165-171.
- Diehl, E. & Diehl-Fleig, Ed. (1997). Primeiro registro de *Zacryptocerus depressus* Klug e de *Z. incertus* Emery (Hymenoptera: Formicidae) no Rio Grande do Sul. *Acta Biologica Leopoldensia*, 19: 225-228.
- Diehl-Fleig, E., Sanhudo, C.E.D. & Diehl-Fleig, Ed. (2000). Mirmecofauna de solo nas dunas da Praia Grande e no Morro da Guarita no município de Torres, RS, Brasil. *Acta Biologica Leopoldensia*, 22: 37-43.
- Diehl, E., Sanhudo, C.E.D. & Diehl-Fleig, Ed. (2004). Ground-dwelling ant fauna of sites with soil high level of copper. *Brazilian Journal of Biology*, 64: 33-39. doi: 10.1590/S1519-69842004000100005
- Diehl, E., Sacchett, F., Albuquerque, E.Z.. (2005a). Riqueza de formigas de solo na Praia da Pedreira, Parque Estadual de Itapuã, Viamão, RS, Brasil. *Revista Brasileira de Entomologia*, 49: 552-556. doi: 10.1590/S0085-56262005000400016
- Diehl, E., Junqueira, L.K. & Berti-Filho, E. (2005b). Ant and termite mound coinhabitants in the wetlands of Santo Antonio da Patrulha, Rio Grande do Sul, Brazil. *Brazilian Journal of Biology*, 65: 431-437. doi: 10.1590/S1519-69842005000300008
- Diehl, E., Florencio, D.F., Schmidt, F.A. & Menzel, L.V.A. (2005c). Riqueza e composição das comunidades de formigas e térmitas na Floresta Nacional de São Francisco de Paula (FLONA-SFP), RS. *Acta Biologica Leopoldensia* 27: 99-106.
- Diehl, E.; Göttert, C.L. & Flores, D.G. (2006). Comunidades de formigas em três espécies utilizadas na arborização urbana em São Leopoldo, Rio Grande do Sul, Brasil. *Bioikos*, 20: 25-32.
- Eggleton, P., Williams, P.M. & Gaston, K.J. (1994). Explaining global termite diversity: productivity or history? *Biodiversity and Conservation* 3: 318-330. doi: 10.1007/BF00056505
- Eggleton, P., Bignell, D.E., Sands, W.A., Waite, B., Wood, T.G. & Lawton, J.H. (1995). The species richness of termites (Isoptera) under differing levels of forest disturbance in the Mbalmayo Forest Reserve, southern Cameroon. *Journal of Tropical Ecology*, 11: 85-98. doi: 10.1017/S0266467400008439
- Eggleton, P. (2000). Global patterns of termite diversity. In T. Abe, D.E. Bignell & M. Higashi (Eds.), (pp. 25-51). *Termites: evolution, sociality, symbioses, ecology*. Dordrecht: Kluwer Academic Publishers
- Eggleton, P. & Tayasu, I. (2001). Feeding groups, lifetypes and the global ecology of termites. *Ecological Research*, 16:

941-960. doi: 10.1046/j.1440-1703.2001.00444.x

Engel, M.S., Grimaldi, D.A. & Krishna, K. (2009). Termites (Isoptera): their phylogeny, classification, and rise to ecological dominance. *American Museum Novitates*, 3650: 1-27.

Evans, T.A., Dawes, T.Z., Ward, P.R. & Lo, N. (2011). Ants and termites increase crop yield in a dry climate. *Natural Communities*, 2: 262 doi: 10.1038/ncomms1257

Fernández, F. (2003). Introducción a las hormigas de la región Neotropical. Bogotá: Instituto de Investigación de Recursos Biológicos Alexander Von Humboldt, 424 p

Florêncio, D.F. & Diehl, E. (2006). Termitofauna (Insecta, Isoptera) em Remanescentes de Floresta Estacional Semidecidual em São Leopoldo, Rio Grande do Sul, Brasil. *Revista Brasileira de Entomologia*, 50: 505-511. doi: 10.1590/S0085-56262006000400011

Flores, D.G., Goettert, C.L., Diehl, E. (2002). Comunidade de formigas em *Inga marginata* (Bignoniaceae) em uma área suburbana. *Acta Biologica Leopoldensia*, 24: 147-155.

Fonseca, R.C. & Diehl, E. (2004). Riqueza de formigas (Hymenoptera, Formicidae) epigéicas em povoamentos de *Eucalyptus* spp. (Myrtaceae) de diferentes idades no Rio Grande do Sul, Brasil. *Revista Brasileira de Entomologia*, 48: 95-100. doi: 10.1590/S0085-56262004000100016

Fontes, L.R. (1995). Sistemática geral de cupins. In E. Berti Filho & L.R. Fontes (Eds.), *Alguns aspectos atuais da biologia e controle de cupins* (pp. 11-17). Piracicaba: FEALQ.

Fontes, L.R. (1998). Novos aditamentos ao "Catálogo dos Isoptera do Novo Mundo", e uma filogenia para os gêneros neotropicais de Nasutitermitinae. In L.R. Fontes & E. Berti Filho (Eds.), *Cupins: o desafio do conhecimento* (pp. 309-412). Piracicaba: FEALQ.

Fontes, L.R. & Araújo, R.L. de. (1999). Os cupins. In F.A.M. Mariconi (Ed.), *Insetos e outros invasores de residências* (pp. 35-90). Piracicaba: FEALQ.

Gathorne-Hardy, F., Syaurani & Eggleton, P. (2001). The effects of altitude and rainfall on the composition of the termites (Isoptera) of the Leuser Ecosystem (Sumatra, Indonesia). *Journal of Tropical Ecology*, 17: 379-393. doi: 10.1017/S0266467401001262

Godoy, M.C.; Laffont, E.R.; Coronel, J.M.; Etcheverry, C. (2012). Termite (Insecta, Isoptera) assemblage of a gallery forest relic from the Chaco province (Argentina): taxonomic and functional groups. *Arxius de Miscellània Zoològica* 10: 55-67.

Gonçalves, C.R. (1961). O gênero *Acromyrmex* no Brasil. *Studia Entomologica*, 4(1-4): 113-180.

Hameister, T.M., Diehl-Fleig, Ed. & Diehl, E. (2003). Comunidades de formigas (Hymenoptera: Formicidae) epigeas no Morro de Itapeva, município de Torres, RS. *Acta Biologica Leopoldensia*, 25: 187-195.

Haubert F., Diehl, E. & Mayhé-Nunes, A. (1998). Mirmecofauna de solo do município de São Leopoldo, RS: Levantamento preliminar. *Acta Biologica Leopoldensia*, 20: 103-108.

Haubert, F. 2006. Riqueza e composição da mirmecofauna de solo no Morro Alto Ferrabraz, município de Sapiranga, RS. São Leopoldo, 2006. 100f. [Dissertação de Mestrado - PPG em Biologia: Diversidade e Manejo de Vida Silvestre/UNISINOS].

Hölldobler, B. & Wilson, E.O. (1990). *The Ants*. Cambridge: Harvard University Press, 732 p

Inoue, T., Takematsu, Y., Yamada, A., Hongoh, Y., Johjima, T., Moriya, S., Sornnuwat, Y., Vongkaluang, C., Ohkuma, M. & Kudo, T. (2006). Diversity and abundance of termites along an altitudinal gradient in Khao Kitchagoot National Park, Thailand. *Journal of Tropical Ecology*, 22: 609-612. doi: 10.1017/S0266467406003403

Jones, D.T. (2000). Termite assemblages in two distinct montane forest types at 1000 m elevation in the Maliau Basin, Sabah. *Journal of Tropical Ecology*, 16: 271-286. doi: 10.1017/S0266467400001401

Jones, D.T. & Eggleton, P. (2000). Sampling termite assemblages in tropical forests: testing a rapid biodiversity assessment protocol. *Journal of Applied Ecology*, 37: 191-203. doi: 10.1046/j.1365-2664.2000.00464.x

Jones, D.T. & Eggleton, P. (2011) Global biogeography of termites: a compilation of sources. In D.E. Bignell, Y. Roisin & N. Lo (Eds.) *Biology of termites: a modern synthesis* (pp. 477-517). Dordrecht: Springer. doi: 10.1007/978-90-481-3977-4_17.

Jones, D.T., Susilo, F.X., Bignell, D.E., Hardiwinotos, S., Gillinson, A.N. & Eggleton, P. (2003). Termite assemblages collapse along a land-use intensification gradient in lowland central Sumatra, Indonesia. *Journal of Applied Ecology*, 40: 380-391. doi: 10.1046/j.1365-2664.2003.00794.x

Jouqueta, P., Traoréc, S., Choosaid, C., Hartmanna, C. & Bignelle, D. (2011). Influence of termites on ecosystem functioning. Ecosystem services provided by termites. *European Journal of Soil Biology*, 47: 215-222. doi: 10.1016/j.ejsobi.2011.05.005

Junqueira, L.K., Diehl, E. & Diehl-Fleig, Ed. (2001). Formigas (Hymenoptera: Formicidae) visitantes de *Ilex paraguariensis* (Aquifoliaceae). *Neotropical Entomology*, 30: 161-164. doi: 10.1590/S1519-566X2001000100024

Kempf, W.W. (1972). Catálogo abreviado das formigas da região Neotropical (Hymenoptera: Formicidae). *Studia Entomologica*, 15(1-4): 3-344.

Krishna, K., Grimaldi, D.A., Krishna, V., Engel, M.S. (2013). *Treatise on the Isoptera of the World: 1. Introduction*. American Museum of Natural History Museum Bulletin, 377: 1-200. doi: 10.1206/377.1

Kusnezov, N. (1957). Numbers of species of ants in faunae of different latitudes. *Evolution*, 11: 298-299.

- Laffont, E.R., Torales, G.J., Arbino, M.O., Godoy, M.C., Porcel, E.A. & Coronel, J.M. (1998). Termites associadas a *Eucalyptus grandis* W.Hill Ex Maiden en el noroeste de la provincia de Corrientes (Argentina). *Revista de Agricultura*, 73: 201-214.
- Lassau, S.A. & Hochuli, D.F. (2004). Effects of habitat complexity on ant assemblages. *Ecography*, 27: 157-164. doi: 10.1111/j.0906-7590.2004.03675.x
- Lavelle, P., Bignell, D., Lepage, M., Wolters, V., Roger, P., Ineson, P., Heal, O.W. & Dhillion, S. (1997). Soil function in a changing world: the role of invertebrate ecosystem engineers. *European Journal of Soil Biology*, 33: 159-193.
- Lobry de Bruyn, L.A. & Conacher, A.J. (1990). The role of termites and ants in soil modification: a review. *Australian Journal of Soil Research*, 28: 55-93. doi: 10.1071/SR9900055
- Lobry de Bruyn, L.A. (1999). Ants as bio-indicators of soil function in rural environments. *Agriculture, Ecosystems and Environment*, 74(1-3): 425-441. doi: 10.1016/S0167-8809-(99)00047-X
- McCain, C.M. (2009). Global analysis of bird elevational diversity. *Global Ecology and Biogeography*, 18: 346-360. doi: 10.1111/j.1466-8238.2008.00443.x
- McCain, C.M. & Grytnes, J.A. (2010). Elevational gradients in species richness. In *Encyclopedia of Life Sciences (ELS)*. Chichester: John Wiley & Sons, Ltd. doi: 10.1002/9780470015902.a0022548
- Machac, A., Janda, M., Dunn, R.R. & Sanders, N.J. (2011). Elevational gradients in phylogenetic structure of ant communities reveal the interplay of biotic and abiotic constraints on diversity. *Ecography*, 34: 364-371. doi: 10.1111/j.1600-0587.2010.06629.x
- Majer, J.D. (1983). Ants: bio-indicators of mine site rehabilitation, land-use, and land conservation. *Environment Management*, 7: 375-383. doi: 10.1007/BF01866920
- Majer J.D. (1996). Ant recolonization of rehabilitated bauxite mines at Trombetas, Pará, Brazil. *Journal of Tropical Ecology*, 12: 257-273. doi 10.1017/S02667400009445.
- Majer, J.D. & Nichols, O.G. (1998). Long-term re-colonization patterns of ants in Western Australian rehabilitated bauxite mines with reference to their use as indicators of restoration success. *Journal of Applied Ecology*, 35: 161-182. doi: 10.1046/j.1365-2664.1998.00286.x
- Marchioretto, A. & Diehl, E. (2006). Distribuição espaciotemporal de uma comunidade de formigas em um remanescente de floresta inundável às margens de um meandro antigo do rio dos Sinos, São Leopoldo, RS. *Acta Biologica Leopoldensia*, 28: 25-32.
- Marques, T. & Schroeder, J.H. (2014). Ant diversity partitioning across spatial scales: Ecological processes and implications for conserving Tropical Dry Forests. *Austral Ecology*, 39: 72-82. doi: 10.1111/aec.12046
- Mathews, A.G.A. (1977). Studies on termites from the Mato Grosso State, Brazil. Rio e Janeiro: Academia Brasileira de Ciências, 267 p.
- Mayhé-Nunes, A.J. & Diehl, E. (1994). Distribuição de *Acromyrmex* (Hymenoptera: Formicidae) no Rio Grande do Sul. *Acta Biologica Leopoldensia*, 16: 115-118.
- McGeoch, M.A. & Chown, S.L. (1998). Scaling up the value of bioindicators. *Trends in Ecology and Evolution*, 13: 46-47.
- Mill, A.E. (1982). Populações de térmitas (Insecta: Isoptera) em quatro habitats no baixo rio Negro. *Acta Amazonica*, 12: 53-60.
- Mora, P., Miambi, E., Jiménez, J.J., Decaëns, T. & Rouland, C. (2005). Functional complement of biogenic structures produced by earthworms, termites and ants in the neotropical savannas. *Soil Biology and Biochemistry*, 37: 1043-1048. doi: 10.1016/j.soilbio.2004.10.019
- Moraes, A.B. & Diehl, E. (2009). Comunidades de formigas em dois ciclos de cultivo de arroz irrigado na planície costeira do Rio Grande do Sul. *Bioikos*, 23: 29-37.
- New, T.R. (1996). Taxonomic focus and quality control in insect surveys for biodiversity conservation. *Australian Journal of Entomology*, 35: 97-106. doi: 10.1111/j.1440-6055.1996.tb01369.x
- Oliveira, P.S. & Del-Claro, K. (2005). Multitrophic interactions in the Brazilian savanna: ant hemipteran systems, associated insect herbivores, and host plant. In D. Burslem, M. Pinard & S. Hartley (Eds.), *Biotic interaction in the Tropics: their role in the maintenance of species diversity* (pp. 414-438). London: Cambridge University Press.
- Pacheco, R. & Vasconcelos, H.L. (2012). Habitat diversity enhances ant diversity in a naturally heterogeneous Brazilian landscape. *Biodiversity and Conservation*, 21: 797-809. doi: 10.1007/s10531-011-0221-y
- Palin, O.F., Eggleton, P., Malhi, Y., Girardin, C.A.J., Rozas-Dávila, A. & Parr, C.L. (2011). Termite diversity along an Amazon-Andes elevation gradient, Peru. *Biotropica*, 43: 100-107. doi: 10.1111/j.1744-7429.2010.00650.x
- Pinheiro, E.R., Duarte, L.S., Diehl, E. & Hartz, S.M. (2010). Edge effects on epigeic ant assemblages in a grassland-forest mosaic in southern Brazil. *Acta Oecologica*, 36: 365-371. doi: 10.1016/j.actao.2010.03.004
- Rahbek, C. (2005). The role of spatial scale and the perception of large-scale species-richness patterns. *Ecology Letters*, 8: 224-239. doi: 10.1111/j.1461-0248.2004.00701.x
- Rambo, B. (1994). A fisionomia do Rio Grande do Sul. São Leopoldo: Editora Unisinos, 473 p
- Roesch, L.F.W., Vieira, F.C.B., Pereira, V.A., Schünemann,

- A.L., Teixeira, I.F., Senna, A.J.T. & Stefenon, V.M. (2009). The Brazilian Pampa: a fragile biome. *Diversity*, 1: 182-198.
- Ribas, C.R., Campos, R.B.F., Schmidt, F.A. & Solar, R.R.C. (2012). Ants as indicators in Brazil: a review with suggestions to improve the use of ants in environmental monitoring programs. *Psyche*, 2012: Article ID 636749. doi: 10.1155/2012/636749.
- Sacchett, F. & Diehl, E. (2004). Comunidades de formigas de solo no Morro da Grota, Parque Estadual de Itapuã, RS. *Acta Biologica Leopoldensia*, 26: 79-92.
- Sacchett, F., Botton, M. & Diehl, E. (2009). Ant species associated with the dispersal of *Eurhizococcus brasiliensis* (Hempel in Wille) (Hemiptera: Margarodidae) in vineyards of the Serra Gaúcha, Rio Grande do Sul, Brazil. *Sociobiology*, 54: 943-954.
- Schmidt, F.A. & Diehl, E. (2008). What is the effect of soil use on ant communities? *Neotropical Entomology*, 37: 381-388. doi: 10.1590/S1519-566X2008000400005
- SEPLAG. (2008). Atlas socioeconômico do Rio Grande do Sul. <http://www.scp.rs.gov.br/atlas/default.asp>. (accessed date: 30 January, 2014).
- Silva, E.G. & Bandeira, A.G. (1999). Abundância e distribuição vertical de cupins (Insecta, Isoptera) em solo de Mata Atlântica, João Pessoa, Paraíba. *Revista Nordestina de Biologia*, 13(1/2): 13-36.
- Silva, R.R. Da & Brandão, C.R.F. (1999). Formigas (Hymenoptera: Formicidae) como indicadoras da qualidade ambiental e da biodiversidade de outros invertebrados terrestres. *Biotemas*, 12: 55-73.
- Silva, R.R., Feitosa, R.S.M. & Eberhardt, F. (2007). Reduced ant diversity along a habitat regeneration gradient in the southern Brazilian Atlantic Forest. *Forest Ecology and Management*, 240(1-3): 61-69. doi: 10.1016/j.foreco.2006.12.002
- Souza, H.B.A., Alves, W.F. & Vasconcellos, A. (2012). Termite assemblages in five semideciduous Atlantic Forest fragments in the northern coastland limit of the biome. *Revista Brasileira de Entomologia*, 56: 67-72. doi: 10.1590/S0085-56262012005000013
- Torales, G.J., Coronel, J.M., Godoy, M.C., Laffont, E.R. & Romero, V.L. (2008). Additions to the taxonomy and distribution of Isoptera from Argentina. *Sociobiology*, 51: 31-48.
- Ulysséa, M.A., Cereto, C.E., Rosumek, F.B., Silva, R.R. & Lopes, B.C. (2011). Updated list of ant species (Hymenoptera, Formicidae) recorded in Santa Catarina State, southern Brazil, with a discussion of research advances and priorities. *Revista Brasileira de Entomologia*, 55: 603-611. doi: 10.1590/S0085-56262011000400018
- Vasconcellos, A., Bandeira, A.G., Moura, F.M.S., Araújo, V.F.P., Gusmão, M.A. & Constantino, R. (2010). Termite assemblages in three habitats under different disturbance regimes in the semi-arid Caatinga of NE Brazil. *Journal of Arid Environments*, 74: 298-302. doi: 10.1016/j.jaridenv.2009.07.007
- Steffen, L.E. (2006). Riqueza e composição das comunidades de formigas em quatro formas de cultivo de erva-mate (*Ilex paraguariensis* St. Hil. 1822) na encosta superior do Nordeste do Rio Grande do Sul. São Leopoldo, 2006. 51f. [Dissertação de Mestrado - PPG em Biologia: Diversidade e Manejo de Vida Silvestre /UNISINOS].
- Ward, P.S. (2000). Broad-scale patterns of diversity in leaf litter ant communities. In D. Agosti, J.D. Majer, L.E. Alonso & T.R. Schulz (Eds.), *Standard methods for measuring and monitoring biodiversity* (pp. 99-121). Washington: Smithsonian Inst. Press.
- Wilson, E.O. (1997). A situação atual da diversidade biológica. In E.O. Wilson & F.M. Peter (Eds.), *Biodiversidade* (pp. 3-24). Rio de Janeiro: Nova Fronteira.

