



## RESEARCH ARTICLE - ANTS

## First Data on the Ants of the Algerian Islands, with a Specific Focus on the Habibas Islands

ANISSA HENINE-MAOUCHE<sup>1</sup>, RIADH MOULAÏ<sup>1</sup>, SIHEM BAKOUR<sup>1</sup>, SOUFYANE KAHLESSENANE<sup>1</sup>, MOULOUD BENABDI<sup>2</sup>, MOHAMMED HAMIMECHE<sup>1</sup>

1 - Laboratoire Zoologie Appliquée et d'Ecophysiologie Animale, Faculté des Sciences de la Nature et de la Vie, Université de Bejaia, Algérie

2 - Bureau d'études – ETS Abyss Environnemental Services, 18, Lot les Rosiers Ouled Fayet, 16094, Alger, Algérie

### Article History

#### Edited by

Evandro Nascimento Silva, UEFS, Brazil  
 Received 27 November 2023  
 Initial acceptance 25 March 2024  
 Final acceptance 28 May 2024  
 Publication date 12 June 2024

#### Keywords

Formicidae, inventory, island environments, Habibas island, Algerian islands.

#### Corresponding author

Anissa Henine-Maouche  
 Laboratoire Zoologie Appliquée et  
 d'Ecophysiologie Animale, Faculté  
 des Sciences de la Nature et de la Vie,  
 Université de Bejaia, Algérie.  
 E-Mail: anissa.maouche@univ-bejaia.dz

### Abstract

The study of the Algerian islands' myrmecofauna was undertaken for the first time, particularly on Habibas Island. The inventory of Habibas Island identified 13 ant species. The island's plant formations, such as the xerophilous shrub group and the xerophilous halophilous shrub group, show a significant correlation with species distribution. Some species, such as *Pheidole pallidula*, are found in all the formations, while others, such as *Aphaenogaster senilis*, are specific to a single formation. The centesimal and occurrence frequencies of the species vary considerably, highlighting the predominance of *Messor capitatus* on the island. The study also revealed significant differences between Algerian islands and islets in terms of ant species richness. *Tetramorium biskrense*, *Pheidole pallidula*, *Messor barbarus*, *Crematogaster scutellaris*, and *Cataglyphis viatica* are present on several islands, while others are specific to a single island, such as *Crematogaster laestrygon* on the Sahel islet. Statistical modellings, such as simple non-linear and multiple regression, did not show any significant correlation between ant richness and factors such as the island's size, its isolation degree, and its floristic richness. However, the latter does seem to have a more or less significant influence on ant distribution. Spearman's correlation test revealed significant similarities between certain islands, while others showed significant differences. These results highlight the importance of the individual characteristics of each island in determining ant composition.

### Introduction

Insular ecosystems are unique and vulnerable environments characterized by exceptional fauna and flora of great importance to biology and biogeography (Mac Arthur & Wilson, 1967; Quézel et al., 1990; Gros-Desormeaux et al., 2016). The theory of island biogeography predicts that islands are home to fewer species than continents, even with the same surface area, due to factors such as isolation, climate, and higher extinction rates (Mac Arthur & Wilson, 1967). This theory also established that the richness of flora and fauna is proportional to the island's size.

However, despite their small size and geographical isolation, island environments are characterized by their exceptional biodiversity, often containing endemic species.

This biological singularity highlights the importance of islands as privileged research grounds for island biogeography due to their unique ecological characteristics (Krebs et al., 2014).

Biodiversity within these fragile ecosystems has been meticulously explored in the northern Mediterranean region, as shown by the research carried out by Vidal et al. (1998) and Bonnet et al. (1999). On the other hand, the situation is very different on the southern coast of the Mediterranean, in particular in Algeria. Although this region has a long coastline of over 1,200 km, the number of islands and islets is relatively few. Studies assessing the faunal diversity of Algerian islands are rare, except a few studies focusing on specific taxonomic groups, such as breeding seabirds (Boukhalfa, 1990; Moulai, 2006; Moulai et al., 2006; Mouret, 2008; Bougaham & Moulai, 2013; Durand, 2013; Bakour & Moulai, 2019),




reptiles and mammals (Mouret, 2008; Moulai et al., 2015) and island flora (Ben Hamiche-Hanifi & Moulai, 2012; Vela et al., 2012; Vela, 2017; Ghermaoui et al., 2016; Bouyahmed & Moulai, 2018). However, one taxon remains relatively under-researched in island environments, namely entomology, and more specifically myrmecology. Only a few investigations have been undertaken in this field, notably by Aissat & Moulai (2016) and Moulai et al. (2016). As far as Formicidae are concerned, the only studies published in Algeria date back to the work carried out by Bernard (1958) on the Habibas islands and the recent study by Henine-Maouche et al. (2019) on the Bejaia islands.

The Habibas islands, located on Algeria's western coastline, are considered to be the largest islands in Algeria, with a total surface area of almost 40 hectares. They are also the only national archipelago and a unique ornithological and botanical conservatory in the western Mediterranean (PIM, 2013). Our study aims to update the pioneering data of Bernard (1958) on the diversity of the Habibas islands's ants, but also to expand the ants's list of some Algerian island systems. Furthermore, in light of the island biogeography theory's premises, we tested the hypothesis that 1) vegetational types with high plant richness increase ant richness and 2) island area and isolation would be critical factors in increasing ant richness in islands.

## Material and Methods

The investigations were carried out on the Habibas islands, which are situated in the Mediterranean Sea off the town of Ain el Karma, in the Oran region of western Algeria. They are located around 14.8 kilometres north-north-east of Cap Figalo (Bachet et al., 2007). The Habibas archipelago is composed of two main islands (Fig 1). One, to the south-west, covers around 30 hectares at an altitude of 105 metres, while the other, to the north-east, covers around 10 hectares at an altitude of 24 metres. This island complex also includes a remarkable intermediate island of around 1 hectare, reaching a height of 30 metres, as well as several other smaller islands on the periphery. The intermediate islet is difficult to access and has limited vegetation, while the other islets are mainly formed of sea rocks (Pons et al., 2013). Since 1990, the archipelago has been listed as a site of interest in the Mediterranean (Bachet et al., 2007) and was recently classified as an "important plant area" (Yahi et al., 2012), belonging to the Betico-Ricean hotspot (Médail & Quézel, 1997).

These islands are formed entirely of volcanic rocks (Bachet et al., 2007). The Habibas Islands' vegetation is mainly composed of species adapted to maritime conditions, such as sea spray, winds and climatic dryness. Atmospheric humidity has a limited impact on vascular plants but has a greater effect



**Fig 1** - Geographical location of the Habibas Islands (Algeria). (© Reda Kerbush)

on lichens (Bachet et al., 2007). In 2006-2007, 97 species or subspecies of spontaneous plants were recorded, 16 of which were new to the island. Of these species, five (*Spergularia pycnorhiza*, *Brassica spinescens*, *Anthemis chrysantha*, *Rostraria balansae* and *Fumaria munbyi*) are endemic to the Algerian, Moroccan-Algerian or Spanish-Algerian coasts, and two varieties (*Sonchus tenerrimus* var. *amicus* and *Asteriscus maritimus* var. *sericeus*) are closely endemic to the archipelago (Pons et al., 2013).

Regarding fauna, the Habibas Islands host a wide diversity of nesting seabirds (Bachet et al., 2007; Mouret, 2008; Durand, 2013). The herpetofauna includes species such as *Tarentola mauritanica*, *Chalcides ocellatus tiligugu*, *Scelarcis perspicillata perspicillata* and *Macroprotodon abubakeri*. However, the main environmental problem on the island is the invasive presence of the black rat, *Rattus rattus*, and the Yellow-legged Gull, *Larus michahellis* (Bachet et al., 2007).

The archipelago of the Habibas Islands enjoys a semi-arid climate characterised by an alternation between long periods of good weather and short periods of inclement weather. Winds, mainly from the north, are often violent and laden with sea spray. Winds from the south, mainly in summer, have a drying effect (PIM, 2013).


This study was carried out on the main island (35°43'26.61 "N 1° 7'51.56 "W) during the period from 23 to 25 May 2023. To do this, we explored the entire island,

taking into account the different plant formations identified during the study (Fig 2), and used two sampling methods: pitfall traps and manual capture. This involves a meticulous inspection of the island in search of ant nests and trails. In each collection area, we placed 10 pots randomly and regularly spaced 10 m apart and then left them in place for 48 hours. Due to the topographical characteristics of each island, we were unable to set up the traps according to the transect or quadrat protocols. The collected specimens were then transported to the Laboratory of Applied Zoology and Animal Ecophysiology (University of Bejaia, Algeria), where they underwent identification. This was based on a variety of reference sources, including the work of Bernard (1968), Cagniant (1968, 1969, 1970, 1997, 2005, 2006, 2009), as well as specialist websites such as [www.antarea.com](http://www.antarea.com) and [www.antweb.com](http://www.antweb.com).

The other island environments surveyed, located on Algeria's western, central and eastern coasts (Fig 3), are represented by islands and islets ranging in size from 40 ha to 0.44 ha. Most of these islands are not very far from the mainland, generally less than 2 km from the shore, except for the Habibas islands which are separated from the mainland by nearly 10 km. The data used in this research comes from recent publications, expert reports (NRP project, 2013), unpublished data and personal sampling results (Table 1). It should be noted that all the prospectors used the same sampling methods, namely the use of pitfall traps and manual capture.

**Table 1** - List of islands and islets studied along the Algerian coast, and source of data used.

Islands/Islets	Surface area (Ha)	Isolation (m)	Floristic richness	Surveyors/year	Reference
Rachgoun (Aïn Témouchent)	25	2327	55	S. Bakour & R. Moulai, 2016	Bakour and Moulai (2019)
Ile Ronde (Oran)	0,64	590	14	S. Bakour & R. Moulai, 2016	Bakour and Moulai (2019)
Grande Habibas (Oran)	34,5	9749	93	S. Bakour, M. Hamimeche, S. Kahlessenane, M. Benabadjji, 2023	-
Ile Plane (Oran)	2,3	6688	8	S. Bakour & R. Moulai, 2016	Bakour and Moulai (2019)
Tipaza (Tipaza)	0,57	150	5	M. Hamimeche & R. Moulai, 2016	Unpublished
Sandja (Alger)	0,27	1042	1	M. Hamimeche, 2015	Unpublished
Agueli (Alger)	1,17	649	6	K. Hamadi & R. Moulai, 2012	NPR Report (2013)
Tigzirt (Tizi-Ouzou)	0,46	150	36	A.Chelli & R. Moulai, 2012	NPR Report (2013)
Pisans (Bejaia)	1,31	1250	52	A. Henine-Maouche, 2016	Unpublished
El Euch (Bejaia)	1,88	120	60		
Garlic Island (Bejaia)	0,46	113	21	Henine-Maouche & R. Moulai, 2016	Henine-Maouche et al. (2019)
Sahel (Bejaia)	0,46	7	44		
Grand Cavallo Island (Jijel)	3,6	950	82		
Petit Cavallo Island(Jijel)	3,9	750	101	L. Aissat & R. Moulai 2009 et 2010	Aissat (2017)
Grand Cavallo Ilet (Jijel)	0,44	50	23		
Tazerout (Jijel)	0,74	154	11	H. Amrane, M. Ihdene, M. Hamimeche & R. Moulai, 2017	Unpublished
Rahbet Teffah (Skikda)	2,5	1000	18	T. Lachouri, L. Mouloudj & R. Moulai, 2016	Unpublished



**Fig 2** - Map of the different plant formations on Habibas Island and distribution of pitfall traps.

### Data analysis

In order to analyse the results obtained from the myrmecofauna's study of island environments, we used ecological indices such as species richness, which provides an overview of the total number of species present, and centesimal frequency and occurrence, which enable us to assess the prevalence and relative distribution of species in these environments. A Principal Component Analysis (PCA) was carried out to examine the distribution concerning the types of plant formation present on Habibas Island. We also undertook a simple and multiple regression study to explore the correlation between ant richness and various environmental factors such as surface area, isolation and floristic richness. To assess similarities in Formicidae composition between the

seventeen islands and islets studied, we applied Spearman's correlation test. All these analyses were performed with XLSTAT Perpetual 2019.2.2.

### Results

#### *Inventory and ecological analysis of Formicidae on Habibas Island*

The sampling of Habibas Island's Formicidae enabled an inventory to be made of the myrmecofauna living in this environment. Our study identified 13 ant species representing 9 genera and 2 subfamilies: Formicinae (*Camponotus* and *Lepisiota*) et Myrmicinae (*Messor*, *Aphaenogaster*, *Crematogaster*, *Pheidole*, *Tetramorium*, and *Temnothorax*). The list and distribution of ants according to the type of plant formation present on the island (Fig 2) are presented in Table 2.

**Table 2** - List of Formicidae found on Habibas Island and distribution by plant formation.

Nf: Nitrophilous formation; Cg: Chasmophyte grassland; Xsg: Xerophilous shrub grouping; Hg: Halophilous grouping; Acg: Anthemis chrysantha grassland; Hxsg: Haloresistant xerophilous shrub grouping.

Species	Plant formation					
	Nf	Cg	Xsg	Hg	Acg	Hxsg
<i>Camponotus foreli</i>	-	-	+	-	-	-
<i>Camponotus micans</i>	-	+	+	-	+	+
<i>Camponotus pexus</i>	-	+	-	-	+	-
<i>Lepisiota frauenfeldi</i>	-	-	-	+	-	+
<i>Aphaenogaster senilis</i>	-	-	+	-	-	-
<i>Crematogaster laetrygon maura</i>	-	-	+	-	-	+
<i>Messor capitatus</i>	-	-	+	+	-	+
<i>Messor sanctus</i>	-	+	+	-	+	-
<i>Pheidole cicatricosa</i>	-	-	+	+	-	+
<i>Pheidole pallidula</i>	+	+	+	+	+	+
<i>Temnothorax nigritus</i>	-	-	+	-	-	-
<i>Tetramorium biskrense</i>	-	+	+	-	+	+
<i>Tetramorium semilaeve</i>	+	-	+	-	-	+
<b>Richness: 13</b>	<b>2 sp</b>	<b>5 sp</b>	<b>11 sp</b>	<b>4 sp</b>	<b>5 sp</b>	<b>8 sp</b>




Fig 3 - Location of the islands and islets studied along the Algerian coast.

Some species were only sampled in a single plant formation, such as *Temnothorax nigrinus*, *Camponotus foreli* and *Aphaenogaster senilis* found in the xerophilous shrub grouping, while *Pheidole pallidula* is the only species that appeared in all the plant formations. The other species were sampled in two, five or eight different formations.

Principal Component Analysis

A Principal Component Analysis (PCA) was carried out to understand the distribution of ants according to plant formations on the Habibas island (Fig 4). The first principal component (F1) explains 43.95% of the total variance, while

the second principal component (F2) explains 30.92%. The xerophilous shrub grouping has the highest contribution (29.280%) with the F1 axis, while the Chasmophyte grassland has the highest contribution (43.66%) with the F2 axis. The PCA shows a positive correlation between the nitrophilous formation and the Chasmophyte grassland ( $R^2 = 0.735$ ) and between the xerophilous shrub grouping and the halophilous grouping ( $R^2 = 0.670$ ).

The species *Messor capitatus* is associated with the halophilous grouping and the xerophilous shrub grouping, while *Tetramorium semilaeve* makes the greatest contribution to the Chasmophyte grassland (73.153%).




Fig 4 - Principal Component Analysis of the ant species' distribution about plant formations on Habibas Island.

### Centesimal frequencies and occurrences of ants from the Habibas Islands

Frequency values vary considerably from one species to another, ranging from 0.92 to 22.94% for centesimal frequencies and from 16.67 to 100% for occurrences (Fig 5). Some ants have high frequencies of occurrence but low centesimal frequencies, such as *Pheidole pallidula*, *Tetramorium biskrense* and *Camponotus micans*. Although *Messor capitatus* has a frequency of occurrence of 50%, it's the most abundant ant on the island with a frequency of 22.93% (Fig 5). It is followed by *Tetramorium semilaeve* (Fc = 16.51% and Fo = 66.66%) and *Pheidole cicatricosa* (Fc = 11% and Fo = 50%).

### Inventory of Formicidae on islands and islets along the Algerian coast

Sampling of Formicidae on the various islands and islets of the Algerian coast has enabled us to produce an inventory of the myrmecofauna living in these environments. Our sampling enabled us to identify a total of 39 ant species representing 14 genera and 3 subfamilies: Dolichoderinae (*Tapinoma*), Formicinae (*Camponotus*, *Cataglyphis*, *Lasius*, *Lepisiota* and *Plagiolepis*) et Myrmicinae (*Messor*, *Aphaenogaster*, *Crematogaster*, *Pheidole*, *Tetramorium*, and *Temnothorax*) (Table 3).

These results suggest a variation in the distribution of ant species between stations. Some species are present on several islands, such as *Tetramorium biskrense*, *Pheidole pallidula*, *Messor barbarus*, *Crematogaster scutellaris* and *Cataglyphis viatica*, while others are only present on one island, such as *Temnothorax nigrinus*, *Lepisiota frauenfeldi*, *Messor capitatus*, *Messor sanctus*, *Tetramorium semilaeve*, *Crematogaster laestrygon maura*, *Aphaenogaster senilis*, *Camponotus micans*, *Camponotus pexus*, *Camponotus foreli* and *Pheidole cicatricosa*, which were only sampled on the

main island of Les Habibas. *Lasius flavus* and *Crematogaster laestrygon* were sampled only on the Sahel Islet and *Messor antennatus* only on the Pisan island.

### Simple non-linear and multiple regression

For this study, we used a statistical modelling approach, in particular a simple non-linear regression curve supported by multiple regression, to explore the relationships between formicidae richness and various environmental factors, such as the island's size, isolation's degree and its floristic richness. This methodology will enable us to quantify the potential links between these variables and analyse how these factors influence ant diversity on the islands and islets studied.

These statistical tests revealed significant correlations between ant richness and floristic richness ( $R^2 = 0.4355$ ) and a relatively significant correlation with the surface area of the island ( $R^2 = 0.1597$ ). On the other hand, the regression between ant density and soils was not significant ( $R^2 = 0.0023$ ) (Fig 6). The results of the multiple regression carried out with a confidence interval of 95% and the variance analysis did not provide significant results concerning the distribution of ants about the island's characteristics (P-value = 0.115). In addition, the coefficient of determination ( $R^2$ ) was 0.356, suggesting that the model explained only a small part of the variation in ant richness (Fig 7). Nevertheless, the standardised residuals/richness indicates that there is a positive correlation between ant richness and the factors studied for the Sahel Islet and, to a lesser extent, for the Grand Cavallo Islet and Ronde Island. However, for Petit Cavallo Island, there was an unexpected negative correlation for the formicids' distribution. Although its surface area is somewhat large (3.9 Ha) and has a good floristic richness (101 species), there are only 5 ants. The main island of Les Habibas has the greatest richness (13) despite being almost 10 km from the coast.





Fig 5 - Distribution of Habibas ant species according to centesimal and occurrence frequencies.

**Table 3.** Inventory of formicids found on all the islands and islets surveyed in Algeria.

1: Rachgoun Island, 2: Ronde Island, 3: Habibas Island, 4: Plane Island, 5: Tipaza Island, 6: Sandja Island, 7: Agueli Island, 8: Tigzirt Island, 9: El-Euch Island, 10: Garlic Islet, 11: Pisans Island, 12: Sahel Islet, 13: Grand-Cavallo Island, 14: Grand-Cavallo Islet, 15: Petit-Cavallo Island, 16: Tazerout Island, 17: Rahbet Teffah Island.

Species/Stations	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
<i>Tapinoma magnum</i>	-	-	-	-	-	-	-	+	-	-	-	+	-	-	-	-	+
<i>Tapinoma simrothi</i>	-	-	-	-	-	-	-	-	+	+	-	+	+	-	+	-	-
<i>Camponotus vagus</i>	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	-	-
<i>Camponotus foreli</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Camponotus gestroi</i>	+	+	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Camponotus micans</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Camponotus pexus</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Camponotus ruber</i>	-	-	-	-	-	-	-	-	+	-	-	+	-	-	-	-	-
<i>Camponotus sp1</i>	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Camponotus sp2</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Colobopsis truncata</i>	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Cataglyphis viatica</i>	-	-	-	-	-	-	-	+	+	+	-	+	+	+	+	-	+
<i>Lasius flavus</i>	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Lasius sp</i>	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Lepisiota frauenfeldi</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Plagiolepis schmitzii</i>	-	-	-	-	+	-	-	-	+	+	+	-	-	-	-	-	-
<i>Aphaenogaster sardoa</i>	-	+	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Aphaenogaster senilis</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Aphaenogaster testaceo-pilosa</i>	-	+	-	-	-	-	+	-	-	+	+	-	+	-	+	-	-
<i>Aphaenogaster sp</i>	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Crematogaster auberti</i>	-	-	-	-	+	-	-	-	-	-	+	-	+	-	+	-	+
<i>Crematogaster laestrygon</i>	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Crematogaster laestrygon maura</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Crematogaster scutellaris</i>	+	+	-	-	-	-	-	+	-	-	+	+	+	+	+	-	-
<i>Messor antennatus</i>	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Messor barbarus</i>	-	+	-	+	-	-	+	+	+	-	-	+	+	+	+	-	-
<i>Messor capitatus</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Messor sanctus</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Messor structor</i>	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Messor picturatus</i>	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Messor sp1</i>	-	-	-	-	-	-	+	+	-	-	-	-	-	-	-	-	-
<i>Messor sp2</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pheidole cicatricosa</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pheidole pallidula</i>	+	+	+	+	-	+	+	-	+	+	+	+	+	-	+	-	-
<i>Temnothorax nigritus</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Tetramorium biskrense</i>	+	+	+	+	-	-	-	-	+	+	+	-	+	-	+	+	+
<i>Tetramorium semilaeve</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Monomorium salomonis</i>	+	+	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Monomorium subopacum</i>	+	-	-	-	+	-	-	-	-	-	-	-	-	-	-	+	-
<b>Richness: 39</b>	<b>8</b>	<b>9</b>	<b>13</b>	<b>5</b>	<b>3</b>	<b>1</b>	<b>5</b>	<b>5</b>	<b>7</b>	<b>6</b>	<b>7</b>	<b>12</b>	<b>9</b>	<b>5</b>	<b>9</b>	<b>2</b>	<b>4</b>




**Fig 6** - Non-linear regression analysis: Effects of surface area, isolation and floristic richness on ants' richness.

### *Similarity between islands and islets*


To compare the similarities in Formicidae composition between the seventeen islands and islets studied, we used Spearman's correlation test (Fig 8). From this matrix, we can see that Grand Cavallo Island and Petit Cavallo Island have a very high similarity (0.996), which means that they share all their species. Relative similarity was observed between El Euch Island and Garlic Islet (0.595), between Round Island and Plane Island (0.570) and between Plane Island

and Grand Cavallo Island (0.570), suggesting similarities between species.

On the other hand, there is no similarity between Habibas Island and El Euch Island on the one hand and Pisans Island on the other (similarity of 0). The same observation is between Ronde Island and the island of Rahbet Teffah. This means that the ants are completely different on these islands and that they have distinct ant populations. These similarity values can be useful for grouping islands and islets according to their common characteristics (Fig 8).



**Fig 7** - Multiple regression analysis: Effects of size, isolation and floristic richness on ant richness.



**Fig 8** - Correlation map of ants similarities between the islands surveyed.

## Discussions

One of the aims of this study was to update the pioneering data of Bernard (1958) on the myrmecofauna of the Habibas islands. Our sampling enabled us to record 13 species of Formicidae, whereas Bernard sampled only 6 species. These were *Messor sanctus*, *Pheidole pallidula*, *Crematogaster auberti*, *Tetramorium pinucum*, *Lepisiota frauenfeldi* and *Camponotus sylvaticus*. Of the 13 species recorded, only *Pheidole pallidula* and *Lepisiota frauenfeldi* are common to both inventories. Our inventory is therefore missing four species. This can be explained by spatial variability: ant populations can vary from one place to another, even on a small scale (possible differences between Bernard's sampling sites (1958) and ours). or temporal variability: the populations of some ant species may vary from year to year due to environmental factors or competition with other species, knowing that in an island environment, the species extinction rate is between 1,000 and 10,000 times higher than it would be naturally on geological timescales (Gros-Desormeaux et al., 2016).

Lack (1976) suggests that the reduction in species richness observed in this study could also be attributed to barriers to colonisation resulting mainly from ecological factors and the specific communities' organisation. In this study's context, the resistance of island communities to colonisation by new species may be contributing to the reduction in the species richness of Formicidae. It should also be noted that island environments are conducive to species capable of expanding their ecological niche (Lack, 1976). This means that they are better able to exploit a wide range of resources and environmental conditions, demonstrating their ability to adapt to different habitats, adjust their diets according to availability, and withstand changing climatic conditions. Specific depletion on the islands varies according

to the group, with a drastic reduction in the number of predators and modified trophic pyramids, characterised by a widening of the base and a narrowing of the tip (Blondel, 1995). Colonisation is favoured by species that are abundant and widespread on the continent, with a better ability to colonise and persist on an island. Species that are generalist and flexible in their choice of habitats are often favoured (Sara and Morand, 2002). However, some species encounter obstacles such as limited dispersal capacity, excessive specialisation or competition with species already present on the island (McArthur & Wilson, 1967).

The principal component analysis (PCA) carried out provided us with interesting information for understanding the distribution of ants according to the plant formations present on Habibas. The analysis suggests that ants react distinctly to the different plant formations on Habibas Island, which may be due to food preferences or specific ecological niches. For example, *Tetramorium semilaeve* was sampled in nitrophilous formations, in the xerophilous shrubby grouping and in the xerophilous haloresistant shrubby grouping. The nitrophilous formation on the island develops on loose soil enriched with nitrates and phosphates from Yellow-legged Gulls (Bachet et al., 2007). It should be noted that this species prefers xerophilous environments (Lebas et al., 2016), particularly grazed environments and matorrals, and coexists with *Pheidole pallidula* (Cagniant, 1997). The latter was found in all the environments sampled, as it is a common species, widespread everywhere (Cagniant, 1968). *Crematogaster laestrygon maura* has been observed exclusively in the two xerophilous shrub groups. Ants of this species build their nests under stones in more or less degraded areas (Cagniant, 2005) and inhabit the rock gardens of the Atlas (Cagniant, 1962). According to the observations of Bachet et al. (2007), the physiognomy of this grouping is similar to low shrubland.

*Lepisiota frauenfeldi* is a xerophilous (Bernard, 1958) and very halophilous (Bernard, 1971) species, so it is no surprise to find it in both halophilous and xerophilous groupings. *Aphaenogaster senilis*, *Camponotus foreli* and *Temnothorax nigrinus* were only sampled in the xerophilous shrub grouping as these species prefer arid and thermophilous environments with nests under stones (Forel, 1894; Lebas et al., (2016). *Aphaenogaster senilis* builds its nests on shallow, sandy ground at low altitudes (Lebas et al., 2016) and it was in this type of habitat that it was sampled.

The values of the centesimal and occurrence frequencies allowed us to note that *Messor capitatus* is the most abundant species on the island because it is a fairly common species whose colonies are very populous, numbering thousands of workers (Lebas et al., 2016). It should be noted that ants of the *Cataglyphis* genus are absent from this survey, even though they are thermophilic, very xerophilic species that prefer dry soils (Bernard, 1958 and Lebas et al., 2016).

The sampling of formicids in various islands and islets of the Algerian coast enabled us to draw up an inventory of the myrmecofauna living in these environments, with a census of 39 ant species. The greatest richness was observed on Habibas Island (Oran) with 13 species and on the Sahel Islet (Bejaia) with 12 species. This similarity is all the more surprising given that the two islands do not share the same characteristics (surface area, distance from the continent and floristic richness). These results align with those found on the island of Djerba in Tunisia, which has 17 species (Bernard, 1971), Galite (Tunisia) with 12 species and Lampedusa (Italy) with 9 species (Bernard, 1958). The lowest species richness was recorded on the island of Tazerourt (2 species) and the island of Sandja (1 species), as was the island of Linosa (Italy) with 3 species and the island of Lampionne (Italy) with a single species (Bernard, 1958).

The myrmecofauna of the islands and islets of the Algerian coast is very poor compared with that of Bagaux Island (France) with 28 species (Berville et al., 2015), Corsica with 83 species, Sicily with 79 species or Sardinia with 64 species (Casevitz-Weulersse, 1991). Of the 39 species recorded, eight belong to the genus *Messor*, a presence consistent with the region (Berberia) which is probably the cradle of the genus (Bernard, 1958). According to Bernard, the absence of *M. sanctus* and *M. capitatus* from the other islands can be attributed to their competition with *M. barbarus*.

The genus *Camponotus* is one of the largest groups of Formicinae with over 900 species, 24 of which are found in North Africa (Bernard, 1968; Cagniant, 1996). Although the species of this genus have a preference for dry, sloping soils (Bernard, 1968), our study identified only 8 species, characterised by low abundance. Although very common in the Mediterranean, the *Tapinoma* genus is rarely found on Algerian islands and islets. This observation can be explained by its intolerance to the humid air of the summer seasons and not to rain, as these species generally thrive in moist clay soils and regularly watered surfaces (Bernard, 1958 and Lebas et al., 2016).

The species *Monomorium salomonis*, which is widespread in Algeria, is known for its preference for warm and dry environments (Cagniant, 1968 and Lebas et al., 2016) and colonises bare ground (Bernard, 1979). However, it's absent from many of the islands and islets along the Algerian coast, except Rachgoun Island, Plane Island and Round Island. This disparity may be linked to the drier climate on the islands of Algeria's west coast. This region belongs to the semi-arid bioclimatic zone. In comparison, the other Algerian islands in the centre and east are sub-humid and humid bioclimatic zones.

The linear regression curve revealed that the factors studied (surface area, isolation and floristic richness) are not responsible for the ant richness of the islands and islets investigated and suggest that other variables not included in the model play a key role in ant richness on these islands. Only the floristic richness has a significant impact on the distribution of Formicidae. Other factors may play a role in the presence or absence of some species, such as food resources (Lomolino, 2000 and Vanderwoude et al., 2014), ecological conditions (strong winds laden with salt spray) (Bernard, 1958 and Berville et al., 2015), the soil's nature (Bernard, 1971) or the diversity of habitats present on the islands (Casevitz-Weulersse, 1991). However, we point out that further studies must cover a survey time longer than ours in this study or even sample ants in different seasons to overcome the lack of collection of ant species with cryptic life histories or varying population density due to seasonality.

Using Spearman's correlation test to assess compositional similarities, only two islands (Grand Cavallo and Petit Cavallo) were found to be very similar, which can be attributed to their relatively close proximity (around 4 km). Other islands have similarities of between 50 and 60%. On the other hand, the other islands show marked differences from one another. In the island biogeography model approach, an increase in the surface area of a territory increases the chances of offering varied habitat conditions, which translates into an increase in the number of species present (Blondel, 1995). However, it's important to note that this model of island biogeography is not universally applicable. The dynamics of island communities vary according to several parameters, including topographical features, biotope diversity, the intrinsic attributes of colonisers, the number of predators and the history of each island (Lomolino, 2000, Gros-Desormeaux et al., 2016). These variations explain the difference in myrmecofauna on all the islands studied.

For example, although *Crematogaster scutellaris* is commonly associated with oak, particularly cork oak, this species is able to build its nests in various tree species such as fruit trees, ornamental trees and various types of pine (Villemant & Fraval, 2002; Cagniant, 2005; Lebas et al., 2016). On our islands, the nests are built mainly on shrubs such as *Pistacia lentiscus* and *Phillyrea angustifolia*. Despite the ecological variability of this species, it is remarkable that *Crematogaster scutellaris* do not occur on Habibas Island, although the island seems to meet its requirements.

## Conclusion

The study highlighted the diversity and complexity of the myrmecofauna of the Algerian islands. The results suggest specific distribution patterns, but also significant variations between islands. The vegetation type influenced community composition, but island features (Area and isolation) don't. This study contributes to a better understanding of the island's myrmecological fauna and the factors that influence the composition of ant communities on the islands. Finally, for a better understanding of Algerian island ants, it would be useful to extend the investigations to other islands, especially those in eastern Algeria. It's also essential to explore in depth the many factors that influence their presence or absence in order to enrich our knowledge of the ecology of island ants.

## Author's Contributions

SB: Investigation.

SK: Investigation.

MB: Investigation.

MH: Investigation.

AH-M: Concetualization, investigation, data curation, formal analysis, writing-initial draft, writing-review & editing.

RM: Investigation, writing-initial draft, writing-review & editing.

## Acknowledgements

We would like to thank the team at Abyss Environmental Services for their logistical support during the naturalist mission to Habibas Island, and Ms Abdi-Hamecha Lidya (University of Algiers) for her help in identifying an ant species.

## References

Aissat, L. & Moulai, R. (2016). Analysis of insect diversity of three islets in the North-East of Algeria. *Journal of the Entomological Research Society*, 18: 1-21.

Bachet, F., Benhadj, S., Bernard, F., Delauge, J., Harmelin, J., Mante, A., Pascal, M., Tillman, M., Vela, E. & Vidal, P. (2007). Réserve des Iles Habibas Notes naturalistes Petites îles de Méditerranée. Pim, 217p.

Bakour, S. & Moulai, R. (2019). Dietary analysis across breeding seasons of Eleonora's Falcon *Falco eleonora* on the western coast of Algeria. *Ostrich*, 90: 63-72. <https://doi.org/10.2989/00306525.2018.1546774>.

Benhamiche-Hanifi, S. & Moulai, R. (2012). Analyse des phytocénoses des systèmes insulaires des régions de Béjaia et de Jijel (Algérie) en présence du Goéland leucophée (*Larus michahellis*). *Revue d'Écologie (La Terre et La Vie)*, 67: 375-397.

Bernard, F. (1958). Les fourmis des îles Pélagie. Comparaison avec d'autres faunes insulaires. *Rivista di Biologia Coloniale*, 16: 67-79.

Bernard, F. (1968). Faune de l'Europe et du bassin méditerranéen, les fourmis (Hymenoptera Formicidae) d'Europe occidentale et septentrionale. Edition Masson. Paris, 280p.

Bernard, F. (1971). Les fourmis de l'île de Djerba (Tunisie). *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, 62: 3-13.

Bernard, F. (1979). Influence des densités végétales sur les fourmis méditerranéennes. *Écologie des insectes sociaux*. *Compte Rendu Colloque Annuel*, Lausanne. 21-29

Berville, L., Passetti, A. & Ponel, P. (2015). Diversité des Formicidae de la réserve intégrale de l'île de Bagaud (Var, France), avant l'éradication de deux taxa invasifs majeurs: *Rattus rattus* et *Carpobrotus spp.* *Scientific Reports of the Port-Cros National Park*, 29: 23-41.

Blondel, J. (1995). *Biogéographie*. Ed. Masson, 297 p.

Bonnet, V., E. Vidal, F. Médail & T. Tatoni (1999). Analyse diachronique des changements floristiques sur un archipel méditerranéen périurbain (Îles du Frioul, Marseille). *Revue d'Écologie (La Terre et La Vie)*, 54: 3-18.

Bougaham, A.F. & Moulai, R. (2013). Observations sur quelques espèces d'oiseaux de la côte à l'ouest de Jijel (Algérie). *Go-South Bulletin*, 10: 76-85.

Boukhalfa, D., 1990. Observations de quelques espèces d'oiseaux de mer nicheurs sur la côte d'Oran (Algérie). *Revue l'Oiseau et RFO*, 3: 248-251.

Bouyahmed, H. & Moulai, R. (2018). Influence of the yellow-legged gull on soil, composition and organisation of plant communities on some small Islands of Algeria. *Zoology and Ecology*, 28: 300-312. <https://doi.org/10.1080/21658005.2018.1527604>.

Cagniant, H. (1962). Étude de quelques fourmis marocaines. Statistique provisoire des Formicidae du Maroc. *Bulletin de la Société d'Histoire naturelle de l'Afrique du Nord*, 53: 83-118.

Cagniant, H. (1968). Liste préliminaire des fourmis forestières d'Algérie, Résultat obtenu de 1963 à 1966. *Bulletin de la Société d'Histoire Naturelle de Toulouse*, 104: 138-147.

Cagniant, H. (1969). Deuxième liste de fourmis d'Algérie, récoltées principalement en forêt (1er partie). *Bulletin de la Société d'Histoire Naturelle de Toulouse*, 105: 405-430.

Cagniant, H. (1970). Deuxième liste de fourmis d'Algérie (Principalement récoltées en forêt). *Bulletin de la Société d'Histoire Naturelle de Toulouse*, 106: 28-40.

Cagniant, H. (1997). Le genre *Tetramorium* au Maroc (Hymenoptera: Formicidae): Cle et catalogue des espèces.). *Annales de la Societe Entomologique de France*, 1: 89-100.

Cagniant, H. (2005). Les *Crematogaster* du Maroc (Hymenoptera: Formicidae): clé de détermination et commentaires. *Orsis*, 20: 7-12.

Cagniant, H. (2006). Liste actualisée des fourmis du Maroc (Hymenoptera: Formicidae). *Myrmecologische Nachrichten*, 8: 193-200.

- Cagniant, H. (2009). Le genre *Cataglyphis* Foerster, 1850 au Maroc. *Orsis*, 24: 41-71.
- Casevitz-Weulersse, J. (1991). Reproduction et développement des sociétés de *Crematogaster scutellaris* (Olivier, 1791) (Hymenoptera: Formicidae). *Annales de la Société Entomologique de France*, 27: 103-111.
- Durand, J.P. (2013). Suivi ornithologique des populations d'oiseaux marins des îles Habibas – Algérie. *Pim*, 13p.
- Forel, A. (1894). Les formicidés de la province d'Oran. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 30: 114: 1-47.
- Ghermaoui, M., Abdellaoui, K. et Moulai R. (2016). Influence du Goéland leucophaea, *Larus michahellis* sur les formations végétales ouvertes du littoral de Rachgoun (Ouest Oranie, Algérie). *Revue d'Écologie (La Terre et La Vie)*, 71: 250-265.
- Gros-desormeaux, J.R., Tupiassu, L. & Bastos, R.Z. (2016). L'île et le vivant revisités dans la théorie de la biogéographie insulaire: les symptômes du syndrome d'insularité. *Revista GeoAmazonia*, 3: 200-210.  
<http://doi.org/10.17551/2358-1778/geoamazonia.v3n5p200-210>
- Henine-Maouche, A., Aissat, L. & Moulai, R. (2019). Diet of *Cataglyphis bicolor* (Hymenoptera: Formicidae) in an insular condition in northeastern Algeria. *Zoology and Ecology*, 29: 123-133. <https://doi.org/10.35513/21658005.2019.2.8>.
- Krebs, E., Affre, L., Aboucaya, A., Allègre, A., Chenot, J., Montegu, C. & Buisson, E. (2015). Résilience de la flore indigène après éradication des Griffes de sorcière (*Carpobrotus* spp.) sur une île méditerranéenne. *Revue d'Écologie (La Terre et La Vie)*, 12: 80-90.
- Lack, D. (1976). *Island biology*. Oxford: Blackwell scientific publications, 419 p.
- Lebas, C., Galkowski, C., Blatrix, R., & Wegnez, P. (2016). Fourmis d'Europe occidentale. Ed. Delachaux et Niestlé, 415p.
- Lomolino, M.V. (2000). Ecology's most general, yet protean pattern: the species-area relationship. *Journal of Biogeography*, 27: 17-26.
- MacArthur, R.H. & Wilson, E.O. (1967). *The theory of island biogeography*. New Jersey: Princeton University Press, 203 p.
- Medail, F. & Quezel, P. (1997). Hot-spots analysis for conservation of plant biodiversity in the Mediterranean Basin. *Annals of the Missouri Botanical Garden*, 84: 112-127.
- Moulai, R. (2006). Bioécologie de l'avifaune terrestre et marine du Parc National de Gouraya (Doctoral dissertation, INA, Algiers), 182p.
- Moulai, R., Sadoul, N. & Doumandji, S., (2006). Effectifs et biologie de la reproduction du Goéland leucophaea *Larus michahellis* dans la région de Bejaïa (Algérie). *Alauda*, 74: 225-234.
- Moulai R., Benhamiche-Hanifi S., Chelli A., Bougaham A.F., Ghermaoui M. & Hamadi K. (2013) Etude, gestion et conservation de la biodiversité des milieux insulaires de quelques localités de la côte Algérienne. Programme National de Recherche (PNR), DGSRT, MERS, INRAA, 113 p.
- Moulai, R., Bougaham, F.A., Aissat, L., Chelli, A., Ghermaoui, M. & Hamadi, K. (2015). Faune vertébrée de quelques milieux insulaires d'Algérie: Diversité et aspects de conservation. In *Proceeding of the International Congress on Estuaries and coastal protected Area* (pp. 183-190).
- Moulai, R., Aissat, L., Chelli, A. & Hamadi, K., (2016). Insectes des milieux insulaires d'Algérie: Etat des lieux et perspectives de conservation. 10th Maghrebien Références bibliographiques 179 Congress on Marine Sciences and the 5th Franco-Maghrebien Congress of Zoology and Ichthyology, Féz (Maroc). From 8th to 10th December 2016.
- Mouret, V. (2008). Mission de terrain Réserve Naturelle des îles Habibas. *Petites îles de Méditerranée*, 8.
- PIM, (2013). Schéma de gestion de l'archipel des Habibas. Ed. Petite îles de Méditerranée, 65 p.
- Pons, E.C., Clariso, I.E., Casademont, M.C., Fraga i Arguimbau, P. & de Menorca, C.I. (2013). Islands and Plants: Preservation and Understanding of Flora on Mediterranean Islands: 2nd Botanical Conference in Menorca. *Proceedings and Abstracts*. Institut Menorquí d'Estudis.
- Quézel, P., Barbero, M., Bonin, G. & Loisel, R. (1990). Recent plant invasions in the circum-Mediterranean region. *Biological Invasions in Europe and the Mediterranean Basin*, 51-60.
- Sara, M. & Morand, S. (2002). Island incidence and mainland population density: mammals from Mediterranean islands. *Diversity and Distributions*, 8: 1-9.
- Vela, E., (2017). Commentaires sur la flore de l'île de Rachgoun (nord-ouest Algérie). *Revue d'Écologie (La Terre et La Vie)*, 72: 258-268.
- Vela, E., Bougaham, A.F & Moulai, R. (2012). Découverte d'*Allium communatum* Guss. (Alliaceae) en Algérie. *Lagacalia*, 32: 283- 296.
- Vidal, E., Médail, F., Taton, T., Roche, P. & Vidal, P. (1998). Impact of gull colonies on the flora of the Riou archipelago (Mediterranean islands of south-east France). *Biological Conservation*, 84: 235-243.
- Villemant, C. & Fraval, A. (2002). Les insectes ennemis du liège. *Insectes*, 125: 25-30.
- Vanderwoude, C., Haynes, D., Richards, E., Quinquis, B., Moverley, D. & Skelton, P.A. (2014). Gestion des impacts de la petite fourmi de feu (*Wasmannia auropunctata*) en Polynésie française. Programme Régional Océanien de l'Environnement (PROE), 54p.
- Yahi, N., Vela, E., Benhouhou, S., De Belair, G. & Gharzouli, R. (2012). Identifying important plants areas (key biodiversity areas for plants) in northern Algeria. *Journal of Threatened Taxa*, 4: 2753-2765.